

Reference #: **909580**

Radiography Date: 10/15/2013

Practice #:

Date Received: 10/18/2013

**Owner:**

 KATHLENE AMRINE  
 1483 SPENCER RD  
 SALKUM, WA 98582  
 UNITED STATES

**PennHIP Member:**

 DR. CHERYL MC DERMOTT  
 TIMBERLAND VETERINARY SERVICES  
 2150 US HWY 12  
 ETHEL, WA 98542  
 UNITED STATES

**ANIMAL**
**GCH SHOWKAYCE UNDER FYRE (COAL)**

Reg. #: WS40150002

CANINE / AKITA

Microchip: 956000008486584

Date of Birth: 2/14/2012 Sex: M Weight: 85 lbs. Age: 20 mo.

Tattoo:


**RESULTS**

LEFT	Distraction Index (DI)	<b>0.37</b>	DI is greater than 0.30 with no radiographic evidence of DJD. There is an increasing risk of developing DJD as the DI increases; low risk when DI is close to 0.30, high risk when DI is close to 0.70 or above.
	Degenerative Joint Disease (DJD)	<b>None</b>	
	Cavitation	<b>No</b>	
	Other Findings	<b>Not Applicable</b>	
RIGHT	Distraction Index (DI)	<b>0.28</b>	DI is less than or equal to 0.30, with no radiographic evidence of DJD.
	Degenerative Joint Disease (DJD)	<b>None</b>	
	Cavitation	<b>No</b>	
	Other Findings	<b>Not Applicable</b>	

Please note that the PennHIP DI is a measure of hip joint laxity, it does not allude to a "passing" or "failing" hip score.

**LAXITY PROFILE RANKING**

The laxity profile ranking is based on the hip with the greater laxity (DI). This interpretation is based on a cross-section of 328 CANINE animals of the AKITA breed. The median DI for this group is 0.58.



The chart above indicates the ranking of your animal's passive hip laxity (DI) in relation to all CANINE animals of the AKITA breed in our database. This result means that 1) your animal's hips are tighter than approximately 90% of this group of animals (alternatively, 10% of the group has tighter hips than your animal), and 2) your animal's hip laxity is in the tighter half of the laxity profile. Breed-specific evaluations are analyzed semi-annually. Consequently, the average laxity and range of laxity for any given group will change over time.

PennHIP does not make specific breeding recommendations. Selection of sire and dam for mating is the decision of the breeder.

**NOTE: As a minimum breeding criterion, we propose that breeding stock be selected from the population of animals having hip laxity in the tighter half of the breed (to the left of the median mark on the graph). Higher selection pressure equates to more rapid expected genetic change per generation.**

By implementing selection based on passive hip laxity, we expect the breed average DI over the years to move toward tighter hip configuration, meaning lower hip dysplasia susceptibility. The PennHIP database permits scientific adjustment of criteria to reflect these shifts; the average laxity and range of laxity for a particular breed will change over time.